

## **Historic, Archive Document**

Do not assume content reflects current scientific knowledge, policies, or practices.







United States  
Department of  
Agriculture  
Agricultural  
Research  
Service

March 1994

# Agricultural Research Service Progress Report

## The Russian Wheat Aphid Sixth Annual Report



Compiled by James A. Webster and Ruth K. Treat  
USDA, ARS, Plant Science and Water Conservation  
Research Laboratory  
Stillwater, Oklahoma

PSWCRL Prog. Rep. 94-001





## Contents

## Page

Introduction	1
Figures	2
Alternate Hosts for Russian Wheat Aphid	5
Host Plant Resistance	7
Small Grain Germplasm Enhancement	9
RWA-Host Plant Interaction	11
Insect Genetics	14
Biosystematics	15
Biological Control	16
ARS Russian Wheat Aphid Publications	26

## Acknowledgment

Figures 1 and 2 were provided by Dr. David R. Porter, ARS PSWCRL, Stillwater, OK.

Figure 3 was provided by Dr. David McNeal, APHIS PPQ, Hyattsville, MD.

### Publication Citation:

J.A. Webster and R.K. Treat (compilers). 1994. The Russian Wheat Aphid, Sixth Annual Report. U.S. Dep. Agric., Agric. Res. Serv., PSWCRL Prog. Rep. 94-001.

### Individual sections may be cited as:

Author(s). 1994. Section title (or Section title: Laboratory name), pp. X-X. *In* J.A. Webster and R.K. Treat (comp.) The Russian wheat aphid, sixth annual report. U.S. Dep. Agric., Agric. Res. Serv., PSWCRL Prog. Rep. 94-001.





## Introduction

Although it has been eight years since the detection of the Russian wheat aphid (RWA) in the United States, its status as a serious economic pest of wheat and barley has not diminished. The latest economic data provided by Legg and Amosson<sup>1</sup> show that estimated yield losses attributed to RWA in the western United States during the 1991-92 crop year were 25,536,000 bushels with an estimated value of \$74,792,000. This is the second highest yield loss since the collection of RWA economic impact data began in 1987, and 24 times greater than bushels lost in the 1990-91 crop year! As pointed out in last year's report, serious RWA outbreaks will continue to occur when conditions are favorable. During the 1991-92 crop year, 838,000 acres were treated with insecticides for RWA at a cost of \$7,598,000. Estimates of cumulative losses for 1987-92 are \$413.1 million in direct loss (control costs, yield loss, and grazing loss) and 437.7 million in indirect loss (ripple effects sustained by the regional economy) – a total loss approaching \$850.8 million. Additional information about the economic impact of the RWA in the western United States is presented in Figures 1 and 2.

Soon after the original 1986 detection of the RWA in Bailey Co., TX, the pest spread rapidly to the major wheat and barley producing areas of the western United States (Fig. 3) and has probably reached all parts of the country where it is likely to be a problem. Most of the serious infestations and economic losses have occurred west of the 100th meridian. Why the RWA has not become a problem east of this line has generated considerable speculation, but scientific facts to explain this phenomenon remain unavailable.

This is the sixth annual report of research progress on the RWA by the Agricultural Research Service of the USDA. As the report indicates, our research is focused on natural control strategies including biological control and the development of RWA-resistant wheat and barley germplasm lines. Several projects fall within these areas and some overlap. For example, we are increasing our knowledge base in areas such as landscape ecology as it relates to aphid populations. We are also learning more about alternate hosts for the RWA as well as alternate aphid hosts for RWA parasitoids. Our overall goal is to develop improved pest management systems for the RWA.


This report is intended as a brief update on the advances in ARS RWA research that have taken place during the past year. Many of our projects are cooperative with organizations including other ARS scientists and locations, as well as scientists from state agricultural experiment stations and USDA-APHIS. These combined efforts provide an excellent model of a cooperative research endeavor and greatly increase the probability of successfully managing the RWA on a regional basis. Space limitations for this report dictate brevity. Additional information about the RWA or specific projects may be obtained directly from the scientists listed in the individual research reports.

<sup>1</sup> Legg, D. and S. Amosson. 1993. Economic impact of the Russian wheat aphid in the United States: 1991-92. Great Plains Agric. Council. Pub. 147.





**Figure 1. Economic Impact of the Russian Wheat Aphid**




Year	1987	1988	1989	1990	1991	1992
Control Cost	17.2	17.0	20.9	15.4	2.5	7.6
Yield Loss	36.6	108.9	71.2	33.4	3.9	74.8
Indirect Loss	48.7	144.8	94.7	44.5	5.3	99.7






Figure 2. Impact of Russian Wheat Aphid on Small Grain







# RUSSIAN WHEAT APHID 1986-1993



Survey data from National Agricultural Pest Information System 7/11/94





## Alternate Hosts for Russian Wheat Aphid

**Mission:** To identify and characterize RWA-resistant germplasm lines that may serve as breeding resources for both cool- and warm-season cereals and turf, range, and conservation grass species.

**S. Dean Kindler, Research Entomologist**  
**Melissa Johnson, Biological Science Technician**  
**USDA, ARS Plant Science Research Laboratory (PSRL)**  
**Wheat & Other Cereal Crops Research Unit**  
**1301 N. Western**  
**Stillwater, OK 74075**  
**405-624-4189, FAX 405-372-1398**

Possible sources of RWA resistance in the perennial Triticeae are reported for the first time. Resistance was related to the newly defined and realigned genera of the perennial Triticeae based on genomic composition. We could not correlate the degree of RWA resistance with the genomic system of classification. However, genera within the tribe Triticeae can be loosely grouped into three categories: 1) moderately resistant: *Leymus* and *Elytrigia*; 2) tolerant to moderately susceptible: *Agropyron*, *Pseudoroegneria*, *Elymus*, and *Pascopyrum*; and 3) susceptible: *Hordeum* and *Thinopyrum*. Within the tribe Triticeae, the genera *Leymus* and *Elytrigia* are the best sources of resistance. However, within each species, extensive variation for resistant types occurs. Introgression of valuable gene(s) for insect resistance has been achieved by interspecific hybridization between the perennial Triticeae species and the cultivated cereals.

RWA resistance identified in several slender wheatgrass, *Agropyron trachycaulum* (Link) Malte = [*Elymus trachycaulus* (Link) Gould ex Shinnery], plant introductions (PIs) may provide genetic variation necessary to breed RWA-resistant wheat cultivars. Seven PIs and the cultivar Pryor slender wheatgrass were considered resistant to RWA. Six PIs (406307, 183009, 440106, 440100, 440102, and 442444) with low damage scores also had low aphid populations and moderate to low leaf-curling scores. All PIs, including the resistant and susceptible entries, had significantly fewer RWAs surviving after 14 days compared with TAM W-101 wheat. All entries tested for antibiosis level were significantly higher for this trait than TAM W-101 wheat. PIs 440100 and 440102 had the highest level of antibiosis, and both has the lowest level of leaf curling. Results of the preference test also indicated that all slender wheatgrass PIs tested were less preferred by RWA in comparison with wheat. Results of these studies indicate that there is a wide genetic diversity in slender wheatgrass. This diversity should be useful in small grain breeding programs.

David Reed (retired) and Dean Kindler surveyed for RWA in Mendoza Province, Argentina, during September 13-18, 1993. We were accompanied by one Chilean and three Argentinian entomologists. Senör Ortego in earlier surveys for RWA in Mendoza Province had identified nine host plants: *Avena sativa*,

*Bromus unioloides*, *Hordeum comosum*, *H. compressum*, *H. distichum*, *H. vulgare*, *Secale cereale*, *Triticum aestivum*, and one unidentified Graminea. While there, we added *Agropyron elongatum* to the list. It was not difficult to find RWA surviving the winter. In general, overwintering RWA populations were highest on *Bromus unioloides* and *Avena sativa*. In the main cereal growing area of Argentina (Buenos Aires Province and surrounding areas), the presence of the RWA, although not extensively surveyed for, has not been detected, although it has been reported to occur just south of the main cereal growing area. It is feared that if found it will upset the balance between parasites and predators and cereal aphids biocontrol. Presently pesticides are not used in this area, but they may be necessary if RWA becomes economically important.

#### **Publications Since Last Report**

Kindler, S.D. 1993. Alternate hosts for Russian wheat aphid, pp. 4-5. In R.L. Burton and J.A. Webster (comp.) The Russian wheat aphid, fifth annual report. U.S. Dep. Agric., Agric. Res. Serv. PSWCL Prog. Rep. 93-009.

Kindler, S.D., K.B. Jensen, and T.L. Springer. 1993. An Overview: Resistance to the Russian wheat aphid (Homoptera: Aphididae) within the perennial Triticeae. J. Econ. Entomol. 85:1609-1618.

Kindler, S.D., and T.L. Springer. 1993. Screening for Russian wheat aphid resistance in *Hordeum* species. PSWCL Rep. 93-006. 29 pp.

Kindler, S.D., T.L. Springer, and R.J. Tyrl. 1993. Screening for Russian wheat aphid resistance in *Agropyron* species. PSWCL Rep. 93-001. 48 pp.

Kindler, S.D., T.L. Springer, and R.J. Tyrl. 1993. Screening for Russian sheat aphid resistance in *Elymus* species. PSWCL Rep. 93-002. 15 pp.

Kindler, S.D., T.L. Springer, and R.J. Tyrl. 1993. Screening for Russian wheat aphid resistance in *Secale* species. PSWCL Rep. 93-003. 4 pp.

Kindler, S.D., T.L. Springer, and R.J. Tyrl. 1993. Screening for Russian wheat aphid resistance in *Stipa* species. PSWCL Rep. 93-004. 12 pp.

Kindler, S.D., T.L. Springer, and R.J. Tyrl. 1993. Screening for Russian wheat aphid resistance in *Bromus* species. PSWCL Rep. 93-005. 32 pp.

#### **Presentations**

Kindler, S.D., and R.W. Hammon. 1993. Comparison of the host suitability of western wheat aphid with Russian wheat aphid. Annual Meetings, Entomological Society of America.



## **Host Plant Resistance**

**Mission:** To identify resistance sources, study the nature of this resistance, and cooperate with the Small Grain Germplasm Enhancement program in the development and release of RWA-resistant small grain germplasms.

**James A. Webster, Research Entomologist**

**Keith Mirkes, Biological Science Technician**

**Chad Webb, Agriculturalist**

**USDA, ARS Plant Science Research Laboratory (PSRL)**

**Wheat & Other Cereal Crops Research Unit**

**1301 N. Western**

**Stillwater, OK 74075**

**405-624-4126, FAX 405-372-1398**

### **Germplasm Evaluation**

To identify new sources of RWA resistance, the systematic germplasm evaluation program has continued. This year, the majority of material evaluated has been wheat; 9,479 wheat accessions were evaluated, and 40 of these accessions were found to be resistant. To date, over 800 wheat lines have been identified as having some level of resistance. The entire ARS National *Hordeum vulgare* collection will have been tested when the current evaluation of 1800 accessions is completed. To date, 38 barley lines have been found to be resistant, and 67 lines were moderately resistant to moderately susceptible. In a cooperative effort, we have conducted RWA resistance tests of advanced material for several barley and wheat breeding programs. Approximately 1200 lines of wheat from an Australian collection have been received for RWA tests.

### **Other**

New techniques were developed for conducting aphid preference/non-preference tests (antixenosis) with individual cut leaves (see Webster et al., 1994). Previous research has indicated that wheat leaf pubescence is not an effective deterrent for the RWA. Recent research has shown that leaf pubescence also has little effect on greenbugs, but is an effective resistance mechanism against the yellow sugarcane aphid.

### **International Cooperation**

Our laboratory co-hosted the 11th Biennial Plant Resistance to Insects Workshop February 28 - March 2, 1994, and several international participants toured the laboratory during the workshop. Since the last report we also hosted a team of agriculturalists from the People's Republic of China; scientists from the Small Grain Centre, Bethlehem, South Africa; Awarsa College of Agriculture, Ethiopia; International Centre of Insect Physiology, Nairobi, Kenya; Cereal Research Institute, Szeged, Hungary; and the Department of Genetics, University of Cordoba, Spain.

#### **Publications Since Last Report**

Webster, J.A. 1993. Host plant resistance, pp. 6-9. *In* R.L. Burton and J.A. Webster (comp.) The Russian wheat aphid, fifth annual report. U.S. Dep. Agric., Agric. Res. Serv., PSWCL Prog. Rep. 93-009.

Webster, J.A., F. Dutoit, and T.W. Popham. 1993. Fecundity comparisons of the Russian wheat aphid (Homoptera: Aphididae) in Bethlehem, South Africa, and in Stillwater, Oklahoma. *J. Econ. Entomol.* 86:544-548.

Webster, J.A., C. Inayatullah, M. Hamissou, and K.A. Mirkes. 1994. Leaf pubescence effects in wheat on yellow sugarcane aphids and greenbugs (Homoptera: Aphididae). *J. Econ. Entomol.* 87:231-240.

Webster, J.A., D.R. Porter, C.A. Baker, and D.W. Mornhinweg. 1993. Russian wheat aphid (Homoptera: Aphididae) resistance in barley: Effects on aphid feeding. *J. Econ. Entomol.* 86: 1603-1608.



## **Small Grain Germplasm Enhancement**

**Mission:** To identify, characterize, and introgress genes conferring RWA resistance for small grain germplasm enhancement.

**David R. Porter, Research Geneticist**

**Cheryl A. Baker, Geneticist**

**Dolores W. Mornhinweg, Geneticist**

**Rita Veal, Biological Science Technician**

**USDA, ARS Plant Science Research Laboratory (PSRL)**

**Wheat & Other Cereal Crops Research Unit**

**1301 N. Western**

**Stillwater, OK**

**405-624-4212, 405-624-4251, FAX 405-372-1398**

RWA-resistant wheat germplasms STARS-9302W and STARS-9303W were released. These hard red facultative germplasms were developed from crosses between an unadapted RWA-resistant parent (PI 149898) and an adapted RWA-susceptible parent. Both of these germplasms were included in the First Uniform Russian Wheat Aphid Field Test (URWAFT); over 5 locations, these two lines were the highest ranked of any of the improved material.

RWA-resistant barley germplasm STARS-9301B was released to breeders in the fall of 1993. This germplasm was selected out of PI 366450. Genetic tests indicate two genes for resistance in this line.

Genetic analysis of control of RWA resistance in wheat and barley is continuing. Parents, F1, backcross, F2, and F2:3 generations are screened and evaluated for segregation ratios. To date, the genetic control of RWA resistance has been determined for six of the resistant wheat lines: PI's 140207 and 225217 have single dominant genes, PI's 149898, 245462, 366515, and 366616 have two genes. Chi-square analysis of the F2 and BC to both parents suggested mutual epistasis between one additive gene which is more expressed, and one dominant gene which is less expressed.

Allelism tests of identified sources of resistance are ongoing; resistant lines are intercrossed, and the F1 is crossed to a susceptible cultivar. Analysis of segregation ratios within the resulting progeny should reveal whether resistance in the two parent lines is controlled by the same or different gene(s).

Cultivar development programs have been initiated: the mission is to incorporate RWA resistance, through a backcrossing program, into wheat and barley of various classifications which are adapted to different areas of the country.

The transfer of RWA resistance genes from the Russian triticales continues. X-ray therapy is being used to induce chromosomal translocations to transfer the resistance gene from PI 386156 (RWA-resistant triticales) to a hexaploid wheat background.

Newly identified sources of RWA resistance continue to be evaluated in the greenhouse for agronomic characteristics and suitability for use in a breeding program.

#### **Publications Since Last Report**

Baker, C.A., D.R. Porter, and J.A. Webster. 1993. Inheritance of Russian wheat aphid resistance in a winter wheat - PI 149898. Agron. Abstr., S. Branch, p. 15.

Baker, C.A., D.R. Porter, and J.A. Webster. 1993. Inheritance of Russian wheat aphid resistance in two spring wheats. Agron. Abstr., p. 81.

Baker, C.A., D.R. Porter, and J.A. Webster. Registration of STARS-9302W and 9303W Russian wheat aphid-resistant wheat germplasms. Crop Sci. (In Press)

Baker, C.A., J.A. Webster, and D.R. Porter. 1992. Characterization of Russian wheat aphid resistance in a hard white spring wheat. Crop Sci. 32:1442-1446.

Mornhinweg, D.W., and D.R. Porter. 1993. Effect of Russian wheat aphid on yield and yield components of barley. Agron. Abstr., S. Branch, p. 14.

Mornhinweg, D.W., D.R. Porter, and J.A. Webster. 1993. Inheritance of RWA resistance in barley germplasm line STARS-9301B. Agron. Abstr. p. 95.

Mornhinweg, D.W., D.R. Porter, and J.A. Webster. Registration of STARS-9301B Russian wheat aphid-resistant barley germplasm. Crop Sci. (In Press)

Porter, D.R., C.A. Baker, and D.W. Mornhinweg. 1993. Small grain germplasm enhancement, pp. 10-13. *In* R.L. Burton and J.A. Webster (comp.) The Russian wheat aphid, fifth annual report. U.S. Dep. Agric., Agric. Res. Serv., PSWCL Prog. Rep. 93-009.

Porter, D.R., C.A. Baker, and J.A. Webster. 1993. Russian wheat aphid-induced protein alterations in spring wheat. Agron. Abstr., S. Branch, p. 10.

Porter, D.R., J.A. Webster, and C.A. Baker. 1993. Detection of resistance to the Russian wheat aphid in hexaploid wheat. Plant Breeding 110: 157-160.



## **RWA-Host Plant Interaction**

**Mission:** To develop a fundamental understanding of the molecular nature of the physiological and biochemical basis of RWA damage to facilitate the development of resistant germplasm derived from both traditional breeding programs and genetic engineering techniques.

Research in support of this mission is broadbased and involves studies both in the laboratory and greenhouse at Stillwater, OK, and in large-scale field plots located at Brookings, SD. Because each unit is studying a different aspect of the interaction phenomena, we have reported their progress separately to assist interested readers who may wish to seek additional specific information directly from the scientists involved.

**Unit Mission:** To characterize plant physiological and biochemical responses to RWA attack in resistant and susceptible germplasm to identify superior metabolic systems, pathways, or individual components critical to genetic resistance.

**John D. Burd, Research Entomologist**

**Helen Belefant-Miller, Research Associate, Oklahoma State University**

**USDA, ARS Plant Science Research Laboratory (PSRL)**

**Wheat & Other Cereal Crops Research Unit**

**1301 N. Western**

**Stillwater, OK 74075**

**405-624-4212, 405-624-4231, FAX 405-372-1398**

#### **Plant Aphid Interactions**

Studies investigating the effect of RWA on carbohydrate physiology in hard red winter wheat (TAM W-101) have shown that aphid infestation greatly reduced total nonstructural carbohydrate accumulation and altered the nonstructural carbohydrate partitioning patterns between root, stem, and leaf tissues. The greatest reductions in nonstructural carbohydrates were observed in root and leaf tissue and resulted primarily from diminished fructan levels. A concomitant decrease of sucrose in the leaves of the infested plants was consistent with the observed whole-plant decline in fructan levels. Related studies measuring transient changes of chlorophyll *a* fluorescence induction kinetics induced by RWA feeding stress indicate that an early event in the damage response involves the inhibition of  $Q_A^-$  reoxidation. Photoinhibition, characterized by changes in  $F_0$ ,  $F_M$ ,  $F_V$ ,  $F_{VM}$ , and  $F_S$ , occurs during the first two hours of RWA feeding.

#### **Publications Since Last Report**

Belefant-Miller, H.L., D.R. Porter, J.D. Burd, and R.L. Burton. 1993. RWA-host plant interaction: Plant Science Research Laboratory, p. 16. *In* R.L. Burton and J.A. Webster (comp.) The Russian wheat aphid, fifth annual report. U.S. Dep. Agric., Agric. Res. Serv., PSWCL Prog. Rep. 93-009.

Burd, J.D., R.L. Burton, and J.A. Webster. 1993. Evaluation of Russian wheat aphid (Homoptera: Aphididae) damage on resistant and susceptible hosts with comparisons of damage ratings to quantitative plant measurements. *J. Econ. Entomol.* 86: 974-980.

Miller, H.L. and D.R. Porter. 1993. Physiological effects of Russian wheat aphids on a resistant and susceptible barley. *Agron. Abstr., S. Branch.* pp. 14-15.



**Unit Mission:** To study the effect of cropping systems on population dynamics of RWA, and determine how cultural practices affect plant stress and efficiency under RWA infestation.

**Walter E. Riedell, Plant Physiologist**  
**Robert W. Kieckhefer, Research Entomologist**  
**David A. Beck, Biological Science Technician**  
**USDA, ARS Northern Grain Insects Research**  
**Laboratory (NGIRL)**  
**Crops and Entomology Research Unit**  
**Rural Route 3, Box 3**  
**Brookings, SD 57006-9401**  
**605-693-5207, 605-693-5215, FAX 605-693-5240**

#### **RWA Control Using Seed and Rescue Treatments**

The effectiveness of a diversity of foliar insecticides for cereal aphid control were tested and compared with the pre-plant seed treatment Gaucho 480 FS on field plots of Arapaho winter wheat in October 1993. All of the foliar rescue treatments decimated RWA and other cereal aphid populations on the plants. There was no re-appearance of aphid populations on the plants 26 days after treatment. Gaucho provided equivalent control to all rates of foliar insecticides even at cold temperatures, demonstrating the efficacy of this seed treatment for longterm RWA control.

#### **Impact of Foliar Feeding by Different Aphid Species on Root Growth in Winter Wheat**

Previous research from this laboratory has shown that improved nitrogen soil fertility level improves grain yield performance independently of the effects of RWA infestation. To further investigate this phenomenon, an experiment was conducted to elucidate the effect of aphid feeding upon root growth in winter wheat plants. A greenhouse study was run in 1992 where seedling wheat plants were infested with greenbugs, RWAs, or bird cherry-oat aphids for a sum total of 300 aphid days (150 aphids for 2 days, 75 aphids for 4 days, 50 aphids for 6 days, or 25 aphids for 12 days). Plants were evaluated for shoot dry weight, shoot chlorophyll content, and root length. Preliminary results indicate that aphid species that cause leaf chlorosis during feeding (greenbug, RWA) are no more damaging, in terms of reducing root growth, than aphids that do not cause leaf chlorosis (bird cherry-oat aphid). A more complete statistical analysis is underway, and manuscript preparation will take place this year.

#### **Publications Since Last Report**

Riedell, W.E., and R.W. Kieckhefer. 1993. Nitrogen fertilizer management and grain yield loss to Russian wheat aphids. *Cereal Res. Commun.* 21:57-59.

Riedell, W.E., and R.W. Kieckhefer. 1993. RWA-host plant interaction: Northern Grains Insects Research Laboratory, p. 17. *In* R.L. Burton and J.A. Webster (comp.) *The Russian wheat aphid, fifth annual report.* U.S. Dep. Agric., Agric. Res. Serv., PSWCL Prog. Rep. 93-009.

## **Insect Genetics**

**Mission:** To conduct national and worldwide biotypic and genetic studies on the RWA and its parasitoids.

**Kevin A. Shufran, Research Entomologist**  
**USDA, ARS Plant Science Research Laboratory (PSRL)**  
**Wheat & Other Cereal Crops Research Unit**  
**1301 N. Western**  
**Stillwater, OK**  
**405-624-4361, FAX 405-372-1398**

Dr. Kevin A. Shufran will join the PSRL staff in 1994 as a post doctoral Research Associate to continue the research in this area initiated by Dr. Gary Puterka. Dr. Shufran received his Ph.D. from Kansas State University in 1992. His dissertation was on "Phenotypic and Genotypic Variation of Kansas Greenbug (Homoptera: Aphididae) Populations." He has written several publications involving aphid genetics and is well qualified for this position.

Final documentation of previous research results by Dr. Puterka and Dr. Richard Roehrdanz at Fargo, ND, is reported below.

### **Publications Since Last Report**

Puterka, G.J., W.C. Black IV, W.M. Steiner, and R.L. Burton. 1993. Genetic variation and phylogenetic relationships among worldwide collections of the Russian wheat aphid, *Diuraphis noxia* (Mordvilko), inferred from allozyme and RAPD-PCR markers. *Heredity* 70:604-618.

Roehrdanz, R., D.K. Reed, and R.L. Burton. 1993. Use of polymerase chain reaction and arbitrary primers to distinguish laboratory-raised colonies of parasitic Hymenoptera. *Biocontrol* 3:199-206.



## **Biosystematics**

**Mission:** To provide identifications and verifications for RWA and its natural enemies.

**Manya B. Stoetzel, Research Entomologist**  
**USDA, ARS Systematic Entomology Laboratory (SEL)**  
**BARC-West, Bldg. 004, Room 6**  
**Beltsville, MD 20705**  
**301-504-5168, FAX 301-504-6482**

Dr. Stoetzel continued to provide identifications and verifications for *Diuraphis noxia* (Mordvilko). No other activities regarding RWA were reported for 1993.

### **Publications Since Last Report**

Stoetzel, M.B. 1993. Biosystematics, p. 19. *In* R.L. Burton and J.A. Webster (comp.) The Russian wheat aphid, fifth annual report. U.S. Dep. Agric., Agric. Res. Serv., PSWCL Prog. Rep. 93-009.

## **Biological Control**

**Mission:** To develop a fundamental understanding of the biological and ecological relationships between cereal aphids and their natural enemies in the agroecosystem landscape to facilitate the establishment and integration of biological control strategies into sustainable integrated pest management systems.

This multifaceted mission involves scientific expertise from several laboratories. Each unit's specific mission and accomplishments toward that mission will be reported as separate subsections.



**Unit Mission:** To develop strategies in the laboratory, greenhouse, and field for maximum utilization of natural enemies (exotic, naturalized, and endemic) in RWA-infested cereals and grasses.

**Norman C. Elliott, Research Biologist**  
**John D. Burd, Research Entomologist**  
**David K. Reed, Research Entomologist, Retired**  
**Wade French, Biological Science Technician**  
**Brian Jones, Biological Science Technician**  
**Tim Johnson, Biological Technician**  
**USDA, ARS Plant Science Research Laboratory (PSRL)**  
**Wheat & Other Cereal Crops Research Unit**  
**1301 N. Western**  
**Stillwater, OK 74075**  
**405-624-4407, 405-624-4244, 405-624-4231,**  
**FAX 405-372-1398**

Progress has been made toward developing standardized methods for sampling coccinellids in wheat. Removal sampling provides accurate estimates of adult coccinellids at all densities, and larval coccinellid populations at low and intermediate densities; but at high densities the efficiency of removal sampling is too low to provide useful estimates for larvae. Based on preliminary analysis of the data, it appears that sweepnet sampling is the most useful method for sampling larval coccinellids in wheat. Adult coccinellids are most efficiently sampled by visual counting. Because larval coccinellids are not always sampled adequately by visual counting, the sweepnet may be the best method for sampling adults and larvae simultaneously. A sequential sampling scheme for estimating the mean number of adult coccinellids per 2-minute count with known average precision was developed.

As part of a continuing effort to develop a comprehensive management system for all economically important aphid pests of small grains in the Great Plains region, we measured the effect of greenbug infestations on yield components of early- and late-planted spring wheat in each of two years. Resulting data were used to develop a model relating yield to cumulative greenbug feeding days (aphid-days). A multiple linear regression model that included different intercepts for each spring wheat planting and a common slope relating yield to aphid-days provided a good fit to the data. Based on the model, a loss of 41 kg of grain per ha is expected for each 100 aphid-days that accumulate per tiller. The effect of greenbug and bird cherry-oat aphid infestations on yield components of Karl 92 winter wheat is currently being investigated in field trials; the studies include autumn, spring, and autumn and spring infestations of both species.

Results of a three-year study of the field population dynamics of the RWA in western Nebraska are currently being analyzed. The data were used to validate a simulation model of RWA population dynamics in winter wheat and CRP grass fields. The RWA

population model will be expanded into a spatial population dynamics model for use in exploring the effects of landscape structure on RWA population dynamics and management.

For two successive years we studied the seasonal occurrence and abundance of aphid species in alfalfa, canola, sorghum, red clover, cotton, millet, cultivated sunflower, vetch, and native grasses, as well as on several plant species common to uncultivated lands in the Southern and Central Plains. We identified 16 aphid species whose seasonal abundance patterns indicate that they might effectively serve as alternate host/prey "bridges" for aphidophagous predators and parasitoids for use at times when RWA are typically absent from the agricultural landscape or present in extremely low numbers.

In laboratory tests we found that *Diaeretiella rapae* and *Aphidius colemani* imported for classical biological control of the RWA parasitized individuals of several of the common aphid species to which they were exposed for 24 h on caged host plants. Our results indicate that several of the 16 aphid species that commonly occur in wheat, in crops often grown adjacent to wheat, or in adjacent non-agricultural lands may serve as alternate hosts of these parasitoids. Our results indicate that it may be possible to increase probabilities of establishment of RWA natural enemies by conducting releases at locations where adjacent habitats supporting alternate hosts occur. Results also provide knowledge of hosts and associated habitats into which RWA parasitoids might be released to attempt establishment at times when the RWA is temporarily absent in a particular geographic area. They also indicate a potentially important role for habitat diversification to improve biological control of the RWA (and other economically important cereal aphids) by increasing the availability and predictability of alternate hosts. Our results provide no evidence to suggest that inability to rapidly adapt to new hosts will limit the value of habitat diversification as an approach for increasing the effectiveness of these parasitoids in biological control of the RWA in the Great Plains. We are currently conducting similar studies with several geographic isolates of *Aphelinus asychis* and *A. sp. nr. varipes* from Asia.

A system has been successfully devised for the incorporation of radio-labeled markers into RWAs, greenbugs, and their natural enemies. This technique has been adapted for use in both field and greenhouse research, and studies are currently underway to determine RWA natural enemy dispersal patterns, host selection, and efficacies (including competition) in multitrophic systems.

Caged releases consistently resulted in better (greater natural enemy population densities) colonization of release sites than open-field releases. In southeastern Colorado in 1992, caged releases in wheat fields resulted in greater population densities of natural enemies in colonization surveys than releases into CRP. During spring and summer of 1993, 1.23 million RWA natural enemies were released at five locations in eastern Colorado. Releases were made in cooperation USDA, APHIS and the



Colorado Department of Agriculture.

We have expanded our capabilities for quantitative study and simulation modeling by establishing a laboratory with Geographic Information System (ERDAS) and Artificial Intelligence capabilities. We are currently in the process of obtaining, organizing, and analyzing remotely sensed spatial data and constructing spatially explicit habitat suitability maps for the RWA in southeastern Colorado.

Research on interactions among cereal hosts, RWA, and natural enemies are continuing. Research on interactions of entomophagous fungi, resistant and susceptible wheat, and RWA is in the final stages. Tritrophic effects on wheat grasses have been published, and work on long-term tritrophic effects and field tritrophic research is continuing.

A visiting scientist from Oral Roberts University, Dr. H.C. Reed, conducted research on olfactory cues to a common RWA parasitoid, *Diaeretiella rapae*, using a wind tunnel and olfactometer. His work has provided a great deal of preliminary information that will lead to further research this year. An in-house laboratory report and two student reports have been written concerning the research.

We determined thermal thresholds for development of exotic *A. colemani* and *D. rapae*. Lower thermal thresholds are similar to that of the RWA, suggesting that low temperatures may not impose a barrier to population growth by these parasitoids.

We assessed the ability of *Cycloneda ancoralis* to successfully complete development on four aphid species (*Aphis gossypii*, *A. helianthi*, *Diuraphis noxia*, and *Lypaphis erysimis*) common in Great Plains agroecosystems. Results indicate that the coccinellid can complete immature development on all four species, but the species differ qualitatively as a food source for the coccinellid. While the ability to survive on a broad range of prey is insufficient to ensure that the coccinellid can establish and contribute to biological control of *D. noxia* and other cereal aphids, it is a desirable characteristic for successful exploitation of this agroecosystem.

Mummy weight, an easily obtained biological parameter often used as a measure of parasitoid robustness/fitness, was shown to be unreliable due to variance among growth stages. This variance can be eliminated with proper experimental protocols.

Field overwintering studies conducted in Oklahoma in 1990 and 1991 indicate that Soviet and Syrian *D. rapae*, South American *A. asychis* and *A. colemani*, and South American *C. ancoralis* and *H. variegata* can overwinter successfully in some winters. These studies are continuing.

#### **Publications Since Last Report**

Elliott, N.C., B.W. French, D.K. Reed, and S.D. Kindler. Temperature effects on immature development of *Aphidius colemani*. Can. Entomol. (In press)

Elliott, N.C., B.W. French, D.K. Reed, S.D. Kindler, and D.C. Arnold. Effects of temperature and aphid host species on the biology of a Syrian population of *Diaeretiella rapae* M'Intosh (Hymenoptera: Aphididae). Canadian Entomol. (In press)

Elliott, N.C. and G.L. Hein. 1993. A "generic" sampling method for aphid pests of small grains. Agron. Abstr., S. Branch. p. 10.

Elliott, N.C., G.L. Hein, and B.M. Shepard. 1993. Sampling arthropod pests of wheat and rice, pp. 627-666. In L. Pedigo and D.G. Buntin (eds) Sampling Methods for Arthropod Pests in Agriculture. CRC Press, Boca Raton, FL.

Elliott, N.C., R.W. Kieckhefer, and W.C. Kauffman. 1993. Effects of an invading coccinellid, *Coccinella septempunctata* L., on the structure of native coccinellid species assemblages in agricultural crops. Bull. Ecol. Soc. Am. 74:226.

Elliott, N.C., S.D. Kindler, D.K. Reed, and B.W. French. Host change by *Aphidius colemani* (Hymenoptera: Aphididae). Great Lakes Entomol. (In press)

Reed, D.K., J.D. Burd, N.C. Elliott, and R.K. Campbell. 1993. Aspects of tritrophic interactions of Russian wheat aphid, pp. 109-113. in R.D. Lumsden and J.L. Vaughn (eds), Pest Management: Biologically Based Technologies. Am. Chem. Soc., Washington, DC. (Beltsville Symposium XVIII, 1993) 435 pp.

Reed, D.K., N.C. Elliott, and J.D. Burd. 1993. Biological control: Plant Science Research Laboratory, pp. 21-26. In R.L. Burton and J.A. Webster (comp.) The Russian wheat aphid, fifth annual report. U.S. Dep. Agric., Agric. Res. Serv., PSWCL Prog. Rep. 93-009.

Reed, D.K., T. Hamissou, J.D. Burd, N.C. Elliott, and S.D. Kindler. 1993. Relationship of an aphid parasitoid's larval growth characteristics and mummy weight. Southwestern Entomol. 18:245-250.

**Unit Mission:** To import, quarantine, test, ship, release, and establish exotic natural enemies (parasites and predators) for use in classical biological control programs for the RWA.

**Lawrence R. Ertle, Entomologist**  
**Paul W. Schaefer, Research Entomologist**  
**Roger Fuester, Research Entomologist**  
**Ken Swan, Biological Science Technician**  
**Susan Barth, Biological Science Technician**  
**USDA, ARS Beneficial Insects Introduction Research (BIIR)**  
**501 South Chapel Street**  
**Newark, DE 19713**  
**302-731-7330, FAX 302-737-6780**

No activities pertaining to RWA were reported for 1993.

**Publications Since Last Report**

Ertle, L.R., P.W. Schaefer, and R. Fuester. 1993. Biological control: Beneficial Insects Introduction Research Laboratory, pp. 27-28. *In* R.L. Burton and J.A. Webster (comp.) The Russian wheat aphid, fifth annual report. U.S. Dep. Agric., Agric. Res. Serv., PSWCL Prog. Rep. 93-009.



**Unit Mission:** To develop fungal pathogens of RWA, devise strategies for the introduction of fungi for RWA control in the field, and provide taxonomic support to other scientists studying RWA pathogens.

**John D. Vandenberg, Research Entomologist**  
**Jennifer McManus, Biological Science Technician**  
**Richard A. Humber, Microbiologist**  
**USDA, ARS Plant Protection Research Unit (PPRU)**  
**U.S. Plant, Soil and Nutrition Laboratory**  
**Tower Road**  
**Ithaca, NY 14853-0331**  
**607-255-0496, FAX 607-255-2459**

Using the newly-completed (late 1992) insect quarantine facility, T.J. Poprawski conducted preliminary screening of eleven species of fungi (45 isolates) against the RWA in laboratory bioassays during early 1993. A study to test tritrophic interactions among aphids, their host plants, and aphid-pathogenic fungi was done in cooperation with ARS researchers in Stillwater, OK. Partial results indicate that fungus activity is more limited when aphids feed on resistant plant varieties. Details and results of these tests may be obtained from T.J. Poprawski at his new address (c/o USDA ARS Biological Pest Control Research, 2413 E. Highway 83, Weslaco, TX 78596.)

In late 1993, a quantitative bioassay method was developed for testing various fungi against the RWA in the laboratory. Surface-sterilized cut leaves of barley are embedded in agar under aseptic conditions. Known-aged adults are placed on these leaves and then exposed to fungus spore showers. Aphids and their offspring are monitored daily for survival or infection. This method minimizes aphid handling while facilitating observation and data collection. Quantitative comparisons among fungi and among aphid populations treated in various ways are currently underway. Factors being evaluated include fungus species and isolate, aphid age, temperature, and host plant variety.

#### **Publications Since Last Report**

Poprawski, T.J., S.P. Wraight, R.A. Humber. 1993. Biological control: Plant Protection Research Unit, pp. 29-33. *In* R.L. Burton and J.A. Webster (comp.) The Russian wheat aphid, fifth annual report. U.S. Dep. Agric., Agric. Res. Serv., PSWCL Prog. Rep. 93-009.

Wraight, S.P., T.J. Poprawski, W.L. Meyer, and F.B. Peairs. 1993. Natural enemies of Russian wheat aphid (Homoptera: Aphididae) and associated cereal aphid species in spring-planted wheat and barley in Colorado. *Environ. Entomol.* 22:1383-1391.

**Unit Mission:** To collect, study, and import natural enemies of the RWA.

**Keith R. Hopper, Research Entomologist**

**Lawrence A. Lacey, Insect Pathologist**

**David J. Kazmer, Research Entomologist**

**FSN Staff: Kim Chen**

**Dominique Coutinot**

**Guy Mercadier**

**Gerhild Kirk**

**Valerie Noel**

**Natalie Ramualde-Serviat**

**Students: Antonio Mesquita**

**Remi Dabire**

**Xavier Fauvregue**

**Angela Nobrega-Farias**

**Visiting Scientist: Peter Mason, Agriculture Canada**

**USDA, ARS European Biological Control Laboratory (EBCL)**

**Montpellier, France**

**c/o American Embassy - Agriculture**

**Unit 21551**

**APO, AE 09777**

**33-67-04-5600 FAX 33-67-04-5620**

We explored and collected in France, Italy, Greece, and Spain. From these collections, we reared and shipped 7,936 predators and parasitoids from 19 cultures to US quarantines and on to cooperators for release.

Isolates of the fungal pathogens *Paecilomyces fumosoroseus*, *Verticillium lecanii*, and *Beauveria bassiana* proved promising in bioassays against *D. noxia*. Studies on the interaction of fungi and *Aphelinus asychis* are also underway.

In a field exclosure experiment in Colorado, *D. noxia* was 12-fold higher with native natural enemies excluded, but releases of predators and parasitoids did not reduce *D. noxia* density. Similar exclosure experiments were conducted by Z. Basky at the Crop Protection and Soil Conservation Station, Szolnok, Hungary, but the results have not yet been analyzed.

Sampling of *D. noxia* and natural enemy population dynamics was continued. There continue to be two population peaks in the Montpellier area, one in the spring and the other in autumn. As of yet, no sexual reproduction has been found. In the field in France, *D. noxia* density was 16-fold higher with natural enemies excluded. A peak of 25% of *D. noxia* exposed in the field were parasitized by six parasitoid species.

In laboratory experiments, *Leucopis* sp. oviposition depended on *D. noxia* density, three times more eggs were laid on rolled leaves than on open leaves, and *Leucopis* sp. consumed on average 11 aphids during larval development.

Odors from *D. noxia*, barley, and *D. noxia* plus barley attracted female *A. asychis* (field-collected and laboratory-reared) no more than pure air. In laboratory experiments, *A. asychis* males searched significantly more on plants marked by the females with a contact pheromone than on unmarked plants. A mathematical model of male mating behavior explained sex ratio observed in the field.

To determine the genetic basis of traits likely to affect field fitness, we measured juvenile survival, development time, sex ratio, handling time, acceptance, walking speed, flight propensity, longevity, fecundity, and morphology for the  $F_1$  progeny of a diallel cross of isofemale lines of *A. asychis* that have been inbred for over 50 generations. Initial analysis indicated no significant differences between lines in walking speed or flight propensity. However, inbred lines walked more slowly and flew less than outbred lines. Further analyses of these data are in progress.

The suitability of using the randomly amplified polymorphic DNA-polymerase chain reaction (RAPD-PCR) technique as a source of genetic markers for natural enemy release experiments and gene flow studies was investigated. A large number of polymorphic markers (32 variants among 142 total bands) were found in six isofemale lines of *A. asychis* started from the same field. Thus, this technique provides abundant markers in a situation where an alternative technique, allozyme electrophoresis, usually fails for many haplodiploid parasitoids. However, heritability studies indicate that artifactual DNA banding patterns are relatively common in diploid females and this, combined with the dominance of RAPD markers, generally limits the utility of RAPD-PCR to haploid males. In addition, RAPD markers generated by the same primer are often strongly linked and consequently gene flow studies should employ a large number of primers to minimize the effects of marker nonindependence/linkage.

The effects of laboratory mass-rearing on selected fitness components of *A. asychis* were examined using replicated "new" ( $F_2$ ) and "old" ( $F_{21}$ ) mass-rearings. Preliminary analyses indicate that parasitoids from the new mass-rearings are smaller than those from the old mass-rearings. When size differences are taken into account, fitness measures including egg load, longevity, flight propensity, and walking speed do not differ significantly between the new and old mass-rearings. Additional experiments are underway to determine if the larger size of parasitoids in the old mass-rearings is a result of adaptation to laboratory conditions.

A project on geographical variation in cold tolerance of *D. noxia* parasitoids was started with Dr. P. Mason, Agriculture Canada.



**Publications Since Last Report**

Hopper, K.R., L.A. Lacey, and D.J. Kazmer. 1993. Biological control: European Biological Control Laboratory, pp. 34-39. *In* R.L. Burton and J.A. Webster (comp.) The Russian wheat aphid, fifth annual report. U.S. Dep. Agric., Agric. Res. Serv., PSWCL Prog. Rep. 93-009.

Hopper, K.R., and R.T. Roush. 1993. Mate finding, dispersal, number released, and the success of biological control introductions. *Ecol. Entomol.* 18:321-331.

Hopper, K.R., R.T. Roush, and W. Powell. 1993. Management of the genetics of biological-control introductions. *Annu. Rev. Entomol.* 38:27-51.

Hopper, K.R. Potential impacts on threatened and endangered insect species in the continental United States from introductions of parasitic Hymenoptera for the control of insect pest. *In*: H. Hokkanen and J.M. Lynch (eds.), *Benefits and Risks of Biological Control*. Cambridge University Press; London. (In Press)

## ARS Russian Wheat Aphid Publications

- Aidara, S. 1992. Comparison of predatory and parasitic action of *Aphelinus asychis* on the Russian wheat aphid. Report for Diplôme d'Etudes Approfondies en Parasitologie. University of Science and Technology of Languedoc, Montpellier, France. 17 pp.
- Baker, C.A., O.G. Merkle, and R.A. Veal. 1989. Interspecific breeding for Russian wheat aphid resistance. Proc. 18th Hard Red Winter Wheat Work. Conf., pp. 6-7.
- Baker, C.A., D.R. Porter, and J.A. Webster. 1992. Inheritance of Russian wheat aphid resistance in a hard red winter wheat. Agron. Abstr. p. 89.
- Baker, C.A., D.R. Porter, and J.A. Webster. 1993. Inheritance of Russian wheat aphid resistance in a winter wheat - PI 149898. Agron. Abstr., S. Branch, p. 15.
- Baker, C.A., D.R. Porter, and J.A. Webster. 1993. Inheritance of Russian wheat aphid resistance in two spring wheats. Agron. Abstr., S. Branch, p. 81.
- Baker, C.A., J.A. Webster, and D.R. Porter. 1991. Characterization of a Russian wheat aphid-resistant hexaploid wheat plant introduction - PI 140207. Agron. Abstr. p. 86.
- Baker, C.A., J.A. Webster, and D.R. Porter. 1992. Characterization of Russian wheat aphid resistance in a hard white spring wheat. Crop Sci. 32:1442-1446.
- Baker, C.A., J.A. Webster, and D.R. Porter. 1992. Mechanisms of Russian wheat aphid resistance: Identification of antibiosis in hexaploid wheat, pp. 90-93. In W.P. Morrison (comp.) Proceedings of the Fifth Russian Wheat Aphid Conference. Great Plains Agric. Council. Pub. 142.
- Belefant-Miller, H., D.R. Porter, J.D. Burd, and R.L. Burton. 1992. RWA-host plant interaction: Plant Science Research Laboratory, p. 14. In R.L. Burton (comp.) The Russian wheat aphid, fourth annual report. U.S. Dep. Agric., Agric. Res. Serv., PSWCL Prog. Res. 92-001.
- Belefant-Miller, H.L., D.R. Porter, J.D. Burd, and R.L. Burton. 1993. RWA-host plant interaction: Plant Science Research Laboratory, p. 16. In R.L. Burton and J.A. Webster (comp.) The Russian wheat aphid, fifth annual report. U.S. Dep. Agric., Agric. Res. Serv., PSWCL Prog. Rep. 93-009.
- Black, W.C. IV, N.M. DuTeau, G.J. Puterka, J.R. Nechols, and J.M. Pettorini. 1992. Use of random amplified polymorphic DNA polymerase chain reaction (RAPD-PCR) to detect DNA polymorphisms in aphids. Bull. Entomol. Res. 82:151-159.

- Bregitzer, P., A. Hang, P.M. Hayes, S.D. Kindler, and D.W. Mornhinweg. 1991. Stable, Russian wheat aphid-resistant hybrids between barley, *Hordeum vulgare* L. and *H. bulbosum* L. Agron. Abstr. p. 88.
- Burd, J.D., and R.L. Burton. 1989. Impact of Russian wheat aphid feeding damage to wheat seedlings. Proc. 18th Hard Red Winter Wheat Work. Conf., p. 13.
- Burd, J.D., and R.L. Burton. 1992. Characterization of plant damage caused by the Russian wheat aphid (Homoptera: Aphididae). J. Econ. Entomol. 85:2017-2022.
- Burd, J.D., R.L. Burton, and J.A. Webster. 1989. Plant damage caused by Russian wheat aphid feeding. Proc. Russian Wheat Aphid Conf., 25-27 October 1989, Albuquerque, New Mexico, N. Mex. State Univ. Coop Ext. Serv., pp. 144-145.
- Burd, J.D., R.L. Burton, and J.A. Webster. 1993. Evaluation of Russian wheat aphid (Homoptera: Aphididae) damage on resistant and susceptible hosts with comparisons of damage ratings to quantitative plant measurements. J. Econ. Entomol. 86: 974-980.
- Burd, J.D., G.J. Puterka, and R.L. Burton. 1990. Biotypic variation in a worldwide collection of the Russian wheat aphid. II. Characterization of damage to wheat, triticale, and barley. In Aphid-Plant Interactions: Populations to Molecules. Int. Aphid Symp. Abstr. TU20.
- Burd, J.D., and G.W. Todd. 1992. Total chlorophyll and chlorophyll fluorescence profiles of Russian wheat aphid resistant and susceptible wheat, pp. 101-106. In W.P. Morrison (comp.) Proceedings of the Fifth Russian Wheat Aphid Conference. Great Plains Agric. Council. Pub. 142.
- Burton, R.L. (compiler) 1988. The Russian wheat aphid, first annual report. U.S. Dep. Agric., Agric. Res. Serv., Stillwater, OK.
- Burton, R.L. (compiler) 1989. The Russian wheat aphid, second annual report. U.S. Dep. Agric., Agric. Res. Serv., Stillwater, OK.
- Burton, R.L. (compiler) 1990. The Russian wheat aphid, third annual report. U.S. Dep. Agric., Agric. Res. Serv., Stillwater, OK.
- Burton, R.L. (compiler). 1992. The Russian wheat aphid, fourth annual report. U.S. Dep. Agric., Agric. Res. Serv., PSWCL Prog. Rep. 92-001.
- Burton, R.L., and J.D. Burd. 1990. RWA-host plant interactions: Plant Science Research Laboratory, p. 3. In R.L. Burton (comp.) The Russian wheat aphid, third annual report. U.S. Dep. Agric., Agric. Res. Serv., Stillwater, OK.



Burton, R.L., D.R. Porter, C.A. Baker, J.A. Webster, J.D. Burd, and G.J. Puterka. 1991. Development of aphid-resistant wheat germplasm, pp. 203-213. *In* D.A. Saunders (ed.) *Wheat for Nontraditional Warm Areas*. Mexico, D.F.: CIMMYT.

Burton, R.L., and J.A. Webster (compilers). 1993. *The Russian wheat aphid, fifth annual report*. U.S. Dep. Agric., Agric. Res. Serv., PSWCL Prog. Rep. 93-009.

Campbell, R.K., D.K. Reed, J.D. Burd, and R.D. Eikenbary. 1992. Russian wheat aphid and drought stress in wheat: Tritrophic interactions with plant resistance and a parasitoid, pp. 224-234. *In* W.P. Morrison (comp.) *Proceedings of the Fifth Russian Wheat Aphid Conference*. Great Plains Agric. Counc. Pub. 142.

Campbell, R.K., D.K. Reed, J.D. Burd, and R.D. Eikenbary. 1992. Russian wheat aphid and drought stresses in wheat: Tritrophic interactions with *Diaeretiella rapae* and plant resistance, pp. 297-298. *In* S.B.J. Menken, J.H. Visser, and P. Harrewijn (eds.) *Proc. 8th Int. Symp. Insect-Plant Relationships*. Kluwer Acad. Publ.

Elliott, N.C. 1990. Simulation modeling, p. 14. *In* R.L. Burton (comp.) *The Russian wheat aphid, third annual report*. U.S. Dep. Agric., Agric. Res. Serv., Stillwater, OK.

Elliott, N.C. 1991. Biological control of the Russian wheat aphid. *In* *Focus on the Russian wheat aphid*. *Proc. Wheat Technol. Conf.*, 27-28 February 1991, Chappell/Kimball, Nebraska, Univ. of Nebr., pp. 3-5.

Elliott, N.C. 1992. Simulation modeling, p. 21. *In* R.L. Burton (comp.) *The Russian wheat aphid, fourth annual report*. U.S. Dep. Agric., Agric. Res. Serv., PSWCL Prog. Rep. 92-001.

Elliott, N.C. and G.L. Hein. 1993. A "generic" sampling method for aphid pests of small grains. *Agron. Abstr.*, S. Branch. p. 10.

Elliott, N.C., G.L. Hein, and B.M. Shepard. 1993. Sampling arthropod pests of wheat and rice, pp. 627-666. *In* L. Pedigo and D.G. Buntin (eds) *Sampling Methods for Arthropod Pests in Agriculture*. CRC Press, Boca Raton, FL.

Elliott, N.C., and R.W. Kieckhefer. 1990. Dynamics of aphidophagous coccinellid assemblages in small grain fields in eastern South Dakota. *Environ. Entomol.* 19:1320-1329.

Elliott, N.C., and R.W. Kieckhefer. 1990. Biological control: Northern Grain Insect Research Laboratory, p. 17. *In* R.L. Burton (comp.) *The Russian wheat aphid, third annual report*. U.S. Dep. Agric., Agric. Res. Serv., Stillwater, OK.

Elliott, N.C., R.W. Kieckhefer, and R.L. Flanders. 1990. A field cage method for evaluation of coccinellid predators of cereal aphids. *In* *Aphid-Plant Interactions: Populations to Molecules*. *Int. Aphid Symp. Abstr.* M15.

Elliott, N.C., R.W. Kieckhefer, and R.L. Flanders. 1991. A field cage method for evaluating coccinellid predators of cereal aphids, p. 259. *In* D.C. Peters, J.A. Webster, and C.S. Chlouber (eds.) *Aphid-Plant Interactions: Populations to Molecules*. Proc. Int. Aphid Symp., Okla. State Univ., MP-132.

Elliott, N.C., R.W. Kieckhefer, and W.C. Kauffman. 1991. Estimating adult coccinellid populations in wheat fields by removal, sweepnet, and visual count sampling. *Can. Entomol.* 123:1322.

Elliott, N.C., R.W. Kieckhefer, and W.C. Kauffman. 1993. Effects of an invading coccinellid, *Coccinella septempunctata* L., on the structure of native coccinellid species assemblages in agricultural crops. *Bull. Ecol. Soc. Am.* 74:226.

Elliott, N.C., D.K. Reed, J.R. Nechols, R.W. Kieckhefer, S.D. Kindler, R.V. Flanders, B.W. French, and D.C. Arnold. 1992. Evaluating Russian wheat aphid parasitoids for establishment potential in the Great Plains, pp. 160-163. *In* W.P. Morrison (comp.) *Proceedings of the Fifth Russian Wheat Aphid Conference*. Great Plains Agric. Council. Pub. 142.

Ertle, L.R., and P.W. Schaefer. 1990. Biological control: Beneficial Insects Research Laboratory, p. 21. *In* R.L. Burton (comp.) *The Russian wheat aphid, third annual report*. U.S. Dep. Agric., Agric. Res. Serv., Stillwater, OK.

Ertle, L.R., and P.W. Schaefer. 1992. Biological control: Beneficial Insects Introduction Research Laboratory, p. 24. *In* R.L. Burton (comp.) *The Russian wheat aphid, fourth annual report*. U.S. Dep. Agric., Agric. Res. Serv., PSWCL Prog. Rep. 92-001.

Ertle, L.R., P.W. Schaefer, and R. Fuester. 1993. Biological control: Beneficial Insects Introduction Research Laboratory, pp. 27-28. *In* R.L. Burton and J.A. Webster (comp.) *The Russian wheat aphid, fifth annual report*. U.S. Dep. Agric., Agric. Res. Serv., PSWCL Prog. Rep. 93-009.

Feng, M.G., R.M. Nowierski, J.B. Johnson, and T.J. Poprawski. 1992. Epizootics caused by entomophthoralean fungi (Zygomycetes: Entomophthorales) in populations of cereal aphids (Hom., Aphididae) in irrigated small grains in southwestern Idaho, USA. *J. Appl. Entomol.* 113:376-390.

Fuller, B.W., R.L. Gibson, D.D. Walgenbach, N.C. Elliott, and N.A. Boetel. 1989. Evaluation of insecticide treatments for control of Russian wheat aphids on winter wheat. *Insectic. & Acaricide Tests* 14:142.

Gellner, J.L., R.W. Kieckhefer, and M.W. Ferguson. 1989. Assessment of aphid damage in seedling wheat by a slant-board-absorbent technique. *Cereal Res. Commun.* 17:149-151.

Gellner, J.W., R.W. Kieckhefer, and B. Moreno. 1990. Effects of pot size and fertility level on assessment of aphid (Homoptera: Aphididae) feeding damage on greenhouse-grown spring wheat. *J. Kans. Entomol. Soc.* 63:187-192.

Gray, M.E., G.L. Hein, D.D. Walgenbach, and N.C. Elliott. 1990. Effects of Russian wheat aphid (Homoptera: Aphididae) on winter and spring wheat infested during different plant growth stages under greenhouse conditions. *J. Econ. Entomol.* 83:2434-2442.

Gray, M.E., G.L. Hein, D.D. Walgenbach, N.C. Elliott, and R.W. Kieckhefer. 1988. Evaluation of planting-time insecticide treatments and foliar applications against the oat-bird cherry aphid, corn leaf aphid, and Russian wheat aphid. *Insectic. & Acaricide Tests* 13:322-323.

Gruber, F., T.J. Poprawski, and E. Rey. 1991. Survey for natural enemies of *Diuraphis noxia* (Mordvilko) in Eurasia. *IOBC\West Palaearctic Regional Section Bull.* 1991/XIV/4:102-109.

Halbert, S., F. Gruber, and I. Kiriak. 1989. Report on an expedition to obtain natural enemies of *Diuraphis noxia* (Mordvilko) in Moldavia and in the Ukraine (USSR). *Univ. of Idaho, Parma*, 16 pp.

Halbert, S., I. Kiriak, and F. Gruber. 1990. Exploration for natural enemies and survey of Russian wheat aphid in the southern USSR. *Proc. 49th Annu. Pacific Northwest Veg. Conf., Sect. V, Cereal Crop Pests*, pp. 22-23.

Halbert, S., I. Kiriak, F. Gruber, T. Poprawski, and L. Elbersen. 1991. Occurrence of sexual morphs of Russian wheat aphid in North America and the Soviet Union, pp. 121-123. *In* G.D. Johnson (comp.) *Proc. Fourth Russian Wheat Aphid Workshop*, 10-12 October 1990, Bozeman, Montana, Mont. State Univ.

Halbert, S.E., and T.J. Poprawski. 1991. Exploration for natural enemies of *Diuraphis noxia* (Mordvilko) in Soviet Central Asia - April/May 1991. *U.S. Dep. Agric., Agric. Res. Serv. (Ithaca, NY) and Univ. Idaho Rep.* 16 pp.

Hein, G.L. M.E. Gray, N.C. Elliott, and D.D. Walgenbach. 1989. Influence of Russian wheat aphid infestation at various plant growth stages on the growth of spring wheat. *Proc. Russian Wheat Aphid Conf.*, 25-27 October 1989, Albuquerque, NM, N. Mex. State Univ. Coop. Ext. Serv., pp. 155-157.

Hopper, K.R. 1990. Insect genetics: European Parasite Laboratory, pp. 12-13. *In* R.L. Burton (comp.) *The Russian wheat aphid, third annual report*. *U.S. Dep. Agric., Agric. Res. Serv., Stillwater, OK.*



Hopper, K.R. 1992. Biological control: European Biological Control Laboratory, pp 25-28. *In* R.L. Burton (comp.) The Russian wheat aphid, fourth annual report. U.S. Dep. Agric., Agric. Res. Serv., PSWCL Prog. Rep 92-001.

Hopper, K.R., L.A. Lacey, and D.J. Kazmer. 1993. Biological control: European Biological Control Laboratory, pp. 34-39. *In* R.L. Burton and J.A. Webster (comp.) The Russian wheat aphid, fifth annual report. U.S. Dep. Agric., Agric. Res. Serv., PSWCL Prog. Rep. 93-009.

Hopper, K.R., T.J. Poprawski, and F. Gruber. 1990. Biological control: European Parasite Laboratory, pp. 19-20. *In* R.L. Burton (comp.) The Russian wheat aphid, third annual report. U.S. Dep. Agric., Agric. Res. Serv., Stillwater, OK.

Hopper, K.R., and R.T. Roush. 1993. Mate finding, dispersal, number released, and the success of biological control introductions. *Ecol. Entomol.* 18:321-331.

Hopper, K.R., R.T. Roush, and W. Powell. 1993. Management of the genetics of biological-control introductions. *Annu. Rev. Entomol.* 38:27-51.

Humber, R.A. 1990. Fungal pathogens of aphids, pp. 45-56. *In* D.C. Peters, J.A. Webster, and C.S. Chlouber (eds.) Aphid-Plant Interactions: Populations to Molecules. *Proc. Int. Aphid Symp.*, Okla. State Univ., MP-132.

Humber, R.A., S. Wraight, and T.J. Poprawski. 1990. Biological control: Plant Protection Research Laboratory, pp. 17-18. *In* R.L. Burton (comp.) The Russian wheat aphid, third annual report. U.S. Dep. Agric., Agric. Res. Serv., Stillwater, OK.

Kieckhefer, R.W., and N.C. Elliott. 1989. Effect of fluctuating temperatures on development of immature Russian wheat aphids (Homoptera: Aphididae) and demographic statistics. *J. Econ. Entomol.* 82:119-122.

Kieckhefer, R.W., and J.L. Gellner. 1992. Yield losses in winter wheat caused by low-density cereal aphid populations. *Agron. J.* 84:180-183.

Kieckhefer, R.W., J.L. Gellner, W.E. Riedell, and C.D. Dybing. 1989. Physiologic response of spring wheat seedlings to cereal aphid feeding. *Agron. Abstr.* p. 115.

Kindler, S.D. 1990. Alternate hosts for Russian wheat aphid, p. 8. *In* R.L. Burton (comp.) The Russian wheat aphid, third annual report. U.S. Dep. Agric., Agric. Res. Serv., Stillwater, OK.

Kindler, S.D. 1992. Alternate hosts for Russian wheat aphid, pp. 8-9. *In* R.L. Burton (comp.) The Russian wheat aphid, fourth annual report. U.S. Dep. Agric., Agric. Res. Serv., PSWCL Prog. Rep. 92-001.

Kindler, S.D. 1993. Alternate hosts for Russian wheat aphid, pp. 4-5. *In* R.L. Burton and J.A. Webster (comp.) The Russian wheat aphid, fifth annual report. U.S. Dep. Agric., Agric. Res. Serv. PSWCL Prog. Rep. 93-009.

Kindler, S.D., J.P. Breen, and T.L. Springer. 1991. Reproduction and damage by Russian wheat aphid, *Diuraphis noxia* (Mordvilko) (Homoptera: Aphididae) as influenced by fungal endophytes and cool-season turfgrasses. *J. Econ. Entomol.* 84:685-692.

Kindler, S.D., J.P. Breen, and T.L. Springer. 1991. Endophytes in fescue species and associated enhancement of Russian wheat aphid resistance, p. 258. *In* D.C. Peters, J.A. Webster, and C.S. Chlouber (eds.) Aphid-Plant Interactions: Populations to Molecules. Proc. Int. Aphid Symp., Okla. State Univ., MP-132.

Kindler, S.D., L.G. Greer, and T.L. Springer. 1992. Feeding behavior of the Russian wheat aphid (Homoptera: Aphididae) on wheat and resistant and susceptible slender wheatgrass. *J. Econ. Entomol.* 85:2012-2016.

Kindler, S.D., K.B. Jensen, and T.L. Springer. 1993. An Overview: Resistance to the Russian wheat aphid (Homoptera: Aphididae) within the perennial Triticeae. *J. Econ. Entomol.* 85:1609-1618.

Kindler, S.D., and T.L. Springer. 1988. Greenhouse study on the host range of the Russian wheat aphid. Proc. 31st North American Alfalfa Imp. Conf., p. 29.

Kindler, S.D., and T.L. Springer. 1988. USDA-ARS progress report of alternate hosts of Russian wheat aphid, pp. 83-91. *In* F.B. Peairs and S.D. Pilcher (eds.) Proceedings, 2nd Russian Wheat Aphid Workshop, 11-12 October 1988, Denver, CO.

Kindler, S.D., and T.L. Springer. 1989. Alternate hosts of Russian wheat aphid (Homoptera: Aphididae). *J. Econ. Entomol.* 82:1358-1362.

Kindler, S.D., and T.L. Springer. 1989. Alternate hosts and selection for resistance to the Russian wheat aphid within grass genera. Proc. 18th Hard Red Winter Wheat Work. Conf., pp. 14-22.

Kindler, S.D., and T.L. Springer. 1989. Progress in insecticide control studies at Stillwater, Oklahoma: Russian wheat aphid protection in winter wheat. Proc. Russian Wheat Aphid Conf., 25-27 October 1989, Albuquerque, NM, N. Mex. State Univ. Coop. Ext. Serv., pp. 71-76.

Kindler, S.D., and T.L. Springer. 1990. Resistance to the Russian wheat aphid in perennial *Triticeae*. Proc. 32nd North American Alfalfa Imp. Conf., p. 49.

Kindler, S.D., and T.L. Springer. 1990. Alternate hosts and selection for pest resistance to the Russian wheat aphid. *In* Aphid-Plant Interactions: Populations to Molecules. Int. Aphid Symp. Abstr. TU17.

Kindler, S.D., and T.L. Springer. 1991. Resistance to Russian wheat aphid in wild *Hordeum* species. *Crop Sci.* 31:94-97.

Kindler, S.D., and T.L. Springer. 1992. Evaluation of resistance to Russian wheat aphid in *Hordeum bulbosum*. *Ann. Appl. Biol.* 120, (Supplement) Tests Agrochem. Cultiv. 13:90-91.

Kindler, S.D., and T.L. Springer. 1992. Identification of Russian wheat aphid resistance in *Agropyron* species, pp. 83-89. *In* W.P. Morrison (comp.) Proceedings of the Fifth Russian Wheat Aphid Conference. Great Plains Agric. Counc. Pub. 142.

Kindler, S.D., and T.L. Springer. 1993. Screening for Russian wheat aphid resistance in *Hordeum* species. PSWCL Rep. 93-006. 29 pp.

Kindler, S.D., T.L. Springer, and J.P. Breen. 1990. Russian wheat aphid reproduction and plant damage caused to cool-season grasses and effect of fungal endophytes on aphid biology. *In* Aphid-Plant Interactions: Populations to Molecules. Int. Aphid Symp. Abstr. TU16.

Kindler, S.D., T.L. Springer, and K.B. Jensen. 1989. Progress in alternate host studies at Stillwater, Oklahoma: Selecting for resistance to the Russian wheat aphid in related grass species. Proc. Russian Wheat Aphid Conf., 25-27 October 1989, Albuquerque, NM, N. Mex. State Univ. Coop. Ext. Serv., pp. 1-3.

Kindler, S.D., T.L. Springer, and K.B. Jensen. 1991. Differential damage to range grass seedlings by Russian wheat aphid, *Diuraphis noxia* (Homoptera: Aphididae). *J. Econ. Entomol.* 84:333-336.

Kindler, S.D., T.L. Springer, and R.J. Tyrl. 1993. Screening for Russian wheat aphid resistance in *Agropyron* species. PSWCL Rep. 93-001. 48 pp.

Kindler, S.D., T.L. Springer, and R.J. Tyrl. 1993. Screening for Russian wheat aphid resistance in *Elymus* species. PSWCL Rep. 93-002. 15 pp.

Kindler, S.D., T.L. Springer, and R.J. Tyrl. 1993. Screening for Russian wheat aphid resistance in *Secale* species. PSWCL Rep. 93-003. 4 pp.

Kindler, S.D., T.L. Springer, and R.J. Tyrl. 1993. Screening for Russian wheat aphid resistance in *Stipa* species. PSWCL Rep. 93-004. 12 pp.



Kindler, S.D., T.L. Springer, and R.J. Tyrl. 1993. Screening for Russian wheat aphid resistance in *Bromus* species. PSWCL Rep. 93-005. 32 pp.

Kiriak, I., F. Gruber, T. Poprawski, S. Halbert, and L. Elbertson. 1990. Occurrence of sexual morphs of Russian wheat aphid, *Diuraphis noxia* (Homoptera: Aphididae), in several locations in the Soviet Union and the northwestern United States. Proc. Entomol. Soc. Wash. 92:544-547.

Kovalev, O.V., T.J. Poprawski, A.V. Stekolshchikov, A.B. Vereshchagina, and S.A. Gandrabur. 1991. *Diuraphis* Aizenberg (Hom., Aphididae): Key to apterous viviparous females, and review of Russian language literature on the natural history of *Diuraphis noxia* (Kurdjumov, 1913). J. Appl. Entomol. 112:425-436.

Marsh, P.M., M.E. Schauff, and J.W. Johnson. 1989. An identification aid for genera of hymenopterous parasitoids both primary and secondary, that might be reared from the Russian wheat aphid. In F. Peairs (ed.) Russian Wheat Aphid News 3(6):7-11.

Miller, H.L. and D.R. Porter. 1993. Physiological effects of Russian wheat aphids on a resistant and susceptible barley. Agron. Abstr., S. Branch, p. 14-15.

Mornhinweg, D.W., and D.R. Porter. 1991. Inheritance of Russian wheat aphid resistance in barley. Agron. Abstr. p. 107. Mornhinweg, D.W., and D.R. Porter. 1992. Genetic control of Russian wheat aphid resistance in barley, p. 98. In W.P. Morrison (comp.) Proceedings of the Fifth Russian Wheat Aphid Conference. Great Plains Agric. Council. Pub. 142.

Mornhinweg, D.W., and D.R. Porter. 1992. Effect of Russian wheat aphid on yield and yield components of barley. Agron. Abstr. p. 108.

Mornhinweg, D.W., and D.R. Porter. 1993. Effect of Russian wheat aphid on yield and yield components of barley. Agron. Abstr., S. Branch, p. 14.

Mornhinweg, D.W., D.R. Porter, and J.A. Webster. 1993. Inheritance of RWA resistance in barley germplasm line STARS-9301B. Agron. Abstr. p. 95.

Mornhinweg, D.W., D.R. Porter, J.A. Webster, S.D. Kindler, and R.L. Burton. 1992. Evaluation of barley collections and enhancing resistance to the Russian wheat aphid in malting barley germplasm. Annu. Prog. Rep. Malt. Barley Res. pp. 141-144.

Poprawski, T.J. 1989. Survey of *Diuraphis noxia*, grasshoppers and natural enemies in the Soviet Central Asian Republic of Kirgizia, 15 August-16 September 1989. U.S. Dep. Agric., Agric. Res. Serv., European Parasite Laboratory Rep. 26 pp.

Poprawski, T.J. 1990. A search for the Russian wheat aphid, *Diuraphis noxia*, and its natural enemies in the Soviet Republics of Kirgizia and Ouzbekistan, 9-31 May 1990. U.S. Dep. Agric., Agric. Res. Serv., European Parasite Laboratory Rep. 17 pp.

Poprawski, T.J., and V.F. Eastop. 1992. Expected and current success of imported entomophagous species for control of the Russian wheat aphid in the United States. Proc. 19th Int. Congr. Entomol., 28 June-4 July 1992, Beijing, China, p. 12.

Poprawski, T.J., and F. Gruber. 1988. Survey of cereal aphids and natural enemies in Yugoslavia, Greece, Bulgaria and Turkey, 9 May-31 May 1988. U.S. Dep. Agric., Agric. Res. Serv., European Parasite Laboratory Rep. 33 pp.

Poprawski, T.J., and F. Gruber. 1988. Survey of *Diuraphis noxia* in Turkey, 14-19 June 1988. U.S. Dep. Agric., Agric. Res. Serv., European Parasite Laboratory Rep. 7 pp.

Poprawski, T.J., and F. Gruber. 1989. Report on a visit to the Soviet-American biological control laboratory in Kishinev, USSR and on a survey of *Diuraphis noxia* and natural enemies in the Moldavian SSR, 25 May-15 June 1989. U.S. Dep. Agric., Agric. Res. Serv., European Parasite Laboratory Rep. 20 pp.

Poprawski, T.J., G. Mercadier, and F. Gruber. 1992. Biology of the Russian wheat aphid, *Diuraphis noxia* (Mordvilko). Mem. Soc. R. Belge Entomol. 35:337-342. [In French]

Poprawski, T.J., G. Mercadier, and S.P. Wraight. 1992. Interactions between *Diuraphis noxia*, *Zoophthora radicans* and *Aphelinus asychis*: Preliminary results of laboratory studies, pp. 180-188. In W.P. Morrison (comp.) Proceedings of the Fifth Russian Wheat Aphid Conference. Great Plains Agric. Council. Pub. 142.

Poprawski, T.J., E. Rey, and F. Gruber. 1992. Natural enemies of the Russian wheat aphid, *Diuraphis noxia* (Mordvilko). Mem. Soc. R. Belge Entomol. 35:343-346. [In French]

Poprawski, T.J., N.L. Underwood, G. Mercadier, and F. Gruber. 1992. *Diuraphis noxia* (Kurdjumov) - A bibliography on the Russian wheat aphid, 1886-1992. U.S. Dep. Agric., Agric. Res. Serv., Ithaca, NY.

Poprawski, T.J., S.P. Wraight, R.A. Humber. 1993. Biological control: Plant Protection Research Unit, pp. 29-33. In R.L. Burton and J.A. Webster (comp.) The Russian wheat aphid, fifth annual report. U.S. Dep. Agric., Agric. Res. Serv., PSWCL Prog. Rep. 93-009.

Poprawski, T.J., S.P. Wright, and S. Peresypkina. 1992. The barley aphid, *Brachycolus noxius* Mordwilko: An edited translation, pp. 34-55. *In* W.P. Morrison (comp.) Proceedings of the Fifth Russian Wheat Aphid Conference. Great Plains Agric. Counc. Pub. 142.

Poprawski, T.J., S.P. Wright, and N.L. Underwood. 1992. Biological control: Plant Protection Research Laboratory, pp. 28-34. *In* R.L. Burton (comp.) The Russian wheat aphid, fourth annual report. U.S. Dep. Agric., Agric. Res. Serv., PSWCL Prog. Rep. 92-001.

Porter, D.R. 1992. Russian wheat aphid-induced protein profile alterations in barley, pp. 99-100. *In* W.P. Morrison (comp.) Proceedings of the Fifth Russian Wheat Aphid Conference. Great Plains Agric. Counc. Pub. 142.

Porter, D.R., C.A. Baker, and D.W. Mornhinweg. 1990. Small grain germplasm enhancement, p. 7. *In* R.L. Burton (comp.) The Russian wheat aphid, third annual report. U.S. Dep. Agric., Agric. Res. Serv., Stillwater, OK.

Porter, D.R., C.A. Baker, and D.W. Mornhinweg. 1992. Small grain germplasm enhancement, pp. 12-13. *In* R.L. Burton (comp.) The Russian wheat aphid, fourth annual report. U.S. Dep. Agric., Agric. Res. Serv., PSWCL Prog. Rep. 92-001.

Porter, D.R., C.A. Baker, and D.W. Mornhinweg. 1993. Small grain germplasm enhancement, pp. 10-13. *In* R.L. Burton and J.A. Webster (comp.) The Russian wheat aphid, fifth annual report. U.S. Dep. Agric., Agric. Res. Serv., PSWCL Prog. Rep. 93-009.

Porter, D.R., C.A. Baker, R.A. Veal, and D. Mornhinweg. 1990. Russian wheat aphid resistance germplasm enhancement. *In* Aphid-Plant Interactions: Populations to Molecules. Int Aphid Symp. Abstr. TU34.

Porter, D.R., C.A. Baker, R.A. Veal, D.W. Mornhinweg, and J.A. Webster. 1991. Russian wheat aphid resistance germplasm enhancement, p. 274. *In* D.C. Peters, J.A. Webster, and C.S. Chlouber (eds.) Aphid-Plant Interactions: Populations to Molecules. Int. Aphid Symp., Okla. State Univ., MP-132.

Porter, D.R., C.A. Baker, and J.A. Webster. 1992. Russian wheat aphid-induced protein alterations in spring wheat. Agron. Abstr. pp. 195-196.

Porter, D.R., C.A. Baker, and J.A. Webster. 1993. Russian wheat aphid-induced protein alterations in spring wheat. Agron. Abstr., S. Branch, p. 10.



Porter, D.R., D.W. Mornhinweg, J.A. Webster, S.D. Kindler, and R.L. Burton. 1991. Increasing Russian wheat aphid resistance genetic diversity in barley. Proc. 28th Barley Imp. Conf., pp. 51-52.

Porter, D.R., J.A. Webster, and C.A. Baker. 1993. Detection of resistance to the Russian wheat aphid in hexaploid wheat. Plant Breeding 110: 157-160.

Porter, D.R., J.A. Webster, S.D. Kindler, and D.W. Mornhinweg. 1991. Evaluation of barley collections and enhancing resistance to the Russian wheat aphid in malting barley germplasm. Annu. Prog. Rep. Malt. Barley Res. pp. 131-134.

Puterka, G.J. 1990. Insect genetics: Plant Science Research Laboratory, pp. 9-10. In R.L. Burton (comp.) The Russian wheat aphid, third annual report. U.S. Dep. Agric., Agric. Res. Serv., Stillwater, OK.

Puterka, G.J., W.C. Black IV, W.M. Steiner, and R.L. Burton. 1993. Genetic variation and phylogenetic relationships among worldwide collections of the Russian wheat aphid, *Diuraphis noxia* (Mordvilko), inferred from allozyme and RAPD-PCR markers. Heredity 70:604-618.

Puterka, G.J., and J.D. Burd. 1992. Insect genetics: Plant Science Research Laboratory, pp. 16-18. In R.L. Burton (comp.) The Russian wheat aphid, fourth annual report. U.S. Dep. Agric., Agric. Res. Serv., PSWCL Prog. Rep. 92-001.

Puterka, G.J., J.D. Burd, and R.L. Burton. 1990. Biotypic variation in a worldwide collection of the Russian wheat aphid. I. Biotypic variation on barley, wheat, and triticale based on damage ratings and population development. In Aphid-Plant Interactions: Populations to Molecules. Int. Aphid Symp. Abstr. TU19.

Puterka, G.J., J.D. Burd, and R.L. Burton. 1992. Biotypic variation in a worldwide collection of Russian wheat aphid (Homoptera: Aphididae). J. Econ. Entomol. 85:1497-1506.

Puterka, G.J., and R.L. Burton. 1989. Progress report on the genetic studies on the Russian wheat aphid and its parasitoids. Proc. Russian Wheat Aphid Conf., 25-27 October 1989, Albuquerque, NM, N. Mex. State Univ. Coop. Ext. Serv., pp. 146-147.

Puterka, G.J., and R.L. Burton. 1990. Aphid genetics in relation to host plant resistance, pp. 59-69. In D.C. Peters, J.A. Webster, and C.S. Chlouber (eds) Aphid-Plant Interactions: Populations to Molecules. Proc. Int. Aphid Symp., Okla. State Univ., MP-132.

Puttler, B., and D.K. Reed. 1988. USDA-ARS progress report on Russian wheat aphid parasites, pp. 27-30. In F.B. Peairs and S.D. Pilcher (eds.) Proceedings, 2nd Russian Wheat Aphid Workshop, 11-12 October 1988., Denver, CO.

Reed, D.K. 1989. USDA-ARS progress report on biological control activities. *Proc. Russian Wheat Aphid Conf.*, 25-27 October 1989, Albuquerque, NM, N. Mex. State Univ. Coop. Ext. Serv., pp. 99-105.

Reed, D.K., J.D. Burd, N.C. Elliott, and R.K. Campbell. 1993. Aspects of tritrophic interactions of Russian wheat aphid, pp. 109-113. *in* R.D. Lumsden and J.L. Vaughn (eds), *Pest Management: Biologically Based Technologies*. Am. Chem. Soc., Washington, D.C. (Beltsville Symposium XVIII, 1993) 435 pp.

Reed, D.K., and N.C. Elliott. 1990. Biological control: Plant Science Research Laboratory, pp. 16-17. *In* R.L. Burton (comp.) *The Russian wheat aphid, third annual report*. U.S. Dep. Agric., Agric. Res. Serv., Stillwater, OK.

Reed, D.K., and N.C. Elliott. 1992. Biological control: Plant Science Research Laboratory, pp. 22-23. *In* R.L. Burton (comp.) *The Russian wheat aphid, fourth annual report*. U.S. Dep. Agric., Agric. Res. Serv., PSWCL Prog. Rep. 92-001.

Reed, D.K., N.C. Elliott, and J.D. Burd. 1993. Biological control: Plant Science Research Laboratory, pp. 21-26. *In* R.L. Burton and J.A. Webster (comp.) *The Russian wheat aphid, fifth annual report*. U.S. Dep. Agric., Agric. Res. Serv., PSWCL Prog. Rep. 93-009.

Reed, D.K., N.C. Elliott, R.V. Flanders, G.L. Hein, M.A. Kerner, G.J. Michels, Jr., and C.B. Walker. 1992. Caged versus uncaged releases of Russian wheat aphid natural enemies in four states in spring, 1991, pp. 164-169. *In* W.P. Morrison (comp.) *Proceedings of the Fifth Russian Wheat Aphid Conference*. Great Plains Agric. Counc. Pub. 142.

Reed, D.K., T. Hamissou, J.D. Burd, N.C. Elliott, and S.D. Kindler. 1993. Relationship of an aphid parasitoid's larval growth characteristics and mummy weight. *Southwestern Entomol.* 18:245-250.

Reed, D.K., S.D. Kindler, and T.L. Springer. 1992. Interactions of Russian wheat aphid, a hymenopterous parasitoid and resistant and susceptible slender wheatgrass. *Entomol. Exp. Appl.* 64:239-246.

Reed, D.K., J.A. Webster, B.G. Jones, and J.D. Burd. 1990. Tritrophic relationships of Russian wheat aphid (Homoptera: Aphididae), its hymenopterous parasitoid, and resistant and susceptible small grains. *In* *Aphid-Plant Interactions: Populations to Molecules*. Int. Aphid Symp. Abstr. M4.

Reed, D.K., J.A. Webster, B.G. Jones, and J.D. Burd. 1991. Tritrophic relationships of Russian wheat aphid, a hymenopterous parasitoid, and resistant and susceptible small grains. *Biol. Contr.* 1:35-41.

Reed, H.C., D.K. Reed, and N.C. Elliott. 1992. Comparative life table statistics of *Diaeretiella rapae* and *Aphidius matricariae* (Hymenoptera: Aphidiidae) on the Russian wheat aphid (Homoptera: Aphididae) in the laboratory, p. 189. *In* W.P. Morrison (comp.) Proceedings of the Fifth Russian Wheat Aphid Conference. Great Plains Agric. Counc. Pub. 142.

Reed, H.C., D.K. Reed, and N.C. Elliott. 1992. Comparative life table statistics of *Diaeretiella rapae* and *Aphidius matricariae* in the Russian wheat aphid. *Southwest. Entomol.* 17(4):307-312.

Riedell, W.E. 1989. Effects of Russian wheat aphid infestation on barley plant response to drought stress. *Physiol. Plant.* 77:587-592.

Riedell, W.E. 1990. Aphid infestation damage in wheat plants. *Plant Physiol.* 90:S310.

Riedell, W.E. 1990. Tolerance of wheat to Russian wheat aphid: Nitrogen fertilization reduces yield loss. *J. Plant Nutr.* 13:579-584.

Riedell, W.E., and R.W. Kieckhefer. 1990. RWA-host plant interaction: Northern Grain Insect Research Laboratory, pp. 3-4. *In* R.L. Burton (comp.) The Russian wheat aphid, third annual report. U.S. Dep. Agric., Agric. Res. Serv., Stillwater, OK.

Riedell, W.E., and R.W. Kieckhefer. 1992. RWA-host plant interaction: Northern Grain Insect Research Laboratory, p. 15. *In* R.L. Burton (comp.) The Russian wheat aphid, fourth annual report. U.S. Dep. Agric., Agric. Res. Serv., PSWCL Prog. Rep. 92-001.

Riedell, W.E., and R.W. Kieckhefer. 1993. RWA-host plant interaction: Northern Grains Insects Research Laboratory, p. 17. *In* R.L. Burton and J.A. Webster (comp.) The Russian wheat aphid, fifth annual report. U.S. Dep. Agric., Agric. Res. Serv., PSWCL Prog. Rep. 93-009.

Riedell, W.E., and R.W. Kieckhefer. 1993. Nitrogen fertilizer management and grain yield loss to Russian wheat aphids. *Cereal Res. Commun.* 21:57-59.

Roehrdanz, R.L. 1992. Application of PCR techniques for identification of parasites and predators, pp. 190-196. *In* W.P. Morrison (comp.) Proceedings of the Fifth Russian Wheat Aphid Conference. Great Plains Agric. Counc. Pub. 142.

Roehrdanz, R.L. 1992. Insect genetics: Biosciences Research Laboratory, pp. 18-19. *In* R.L. Burton (comp.) The Russian wheat aphid, fourth annual report. U.S. Dep. Agric., Agric. Res. Serv., PSWCL Prog. Rep. 92-001.



Roehrdanz, R., D.K. Reed, and R.L. Burton. 1993. Use of polymerase chain reaction and arbitrary primers to distinguish laboratory-raised colonies of parasitic Hymenoptera. *Biocontrol* 3:199-206.

Springer, T.L., and S.D. Kindler. 1990. Endophyte-enhanced resistance to the Russian wheat aphid and the incidence of endophytes in fescue species. *Proc. Int. Symp. Acremonium/Grass Interactions*, pp. 194-195.

Springer, T.L., S.D. Kindler, T.L. Harvey, and P.W. Stahlman. 1992. Susceptibility of brome grass to the Russian wheat aphid (Homoptera: Aphididae). *J. Econ. Entomol.* 85:1731-1735.

Steiner, W.W.M. 1990. Insect genetics: Biological Control of Insects Research Laboratory, pp. 11-12. *In* R.L. Burton (comp.) The Russian wheat aphid, third annual report. U.S. Dep. Agric., Agric. Res. Serv., Stillwater, OK.

Stoetzel, M.B. 1987. Information on and identification of *Diuraphis noxia* (Homoptera: Aphididae) and other aphid species colonizing leaves of wheat and barley in the United States. *J. Econ. Entomol.* 80:696-704.

Stoetzel, M.B. 1990. The North American species in and related to *Diuraphis* Aizenberg. *Acta Phytopathol. Entomol. Hung.* 25:229-234.

Stoetzel, M.B. 1990. Biosystematics, p. 15. *In* R.L. Burton (comp.) The Russian wheat aphid, third annual report. U.S. Dep. Agric., Agric. Res. Serv., Stillwater, OK.

Stoetzel, M.B. 1990. Some aphids of importance to the southeastern United States (Homoptera: Aphididae). *Fla. Entomol.* 73:580-586.

Stoetzel, M.B. 1992. Biosystematics, p. 20. *In* R.L. Burton (comp.) The Russian wheat aphid, fourth annual report. U.S. Dep. Agric., Agric. Res. Serv., PSWCL Prog. Rep. 92-001.

Stoetzel, M.B. 1992. Taxonomy of the Russian wheat aphid and related species. *Proc. 19th Int. Congr. Entomol.*, 28 June-4 July 1992, Beijing, China, p. 13.

Stoetzel, M.B. 1993. Biosystematics, p. 19. *In* R.L. Burton and J.A. Webster (comp.) The Russian wheat aphid, fifth annual report. U.S. Dep. Agric., Agric. Res. Serv., PSWCL Prog. Rep. 93-009.

Stoetzel, M.B., and R.W. Hammon. 1992. New collections of sexuales of *Diuraphis* (Homoptera: Aphididae) in North America. *Proc. Entomol. Soc. Wash.* 94:598-599.

Webster, J.A. 1989. Current status of the Russian wheat aphid and progress in plant resistance research. Proc. 18th Hard Red Winter Wheat Work. Conf., pp. 305.

Webster, J.A. 1990. Resistance in triticale to the Russian wheat aphid (Homoptera: Aphididae). J. Econ. Entomol. 83:1091-1095.

Webster, J.A. 1990. Host plant resistance, pp. 5-6. In R.L. Burton (comp.) The Russian wheat aphid, third annual report. U.S. Dep. Agric., Agric. Res. Serv., Stillwater, OK.

Webster, J.A. 1991. Developing aphid-resistant cultivars, pp. 87-99. In D.C. Peters, J.A. Webster, and C.S. Chlouber (eds.) Aphid-Plant Interactions: Populations to Molecules. Proc. Int. Aphid Symp., Okla. State Univ., MP-132.

Webster, J.A. 1992. Host plant resistance, pp. 10-11. In R.L. Burton (comp.) The Russian wheat aphid, fourth annual report. U.S. Dep. Agric., Agric. Res. Serv., PSWCL Prog. Rep. 92-001.

Webster, J.A. 1993. Host plant resistance, pp. 6-9. In R.L. Burton and J.A. Webster (comp.) The Russian wheat aphid, fifth annual report. U.S. Dep. Agric., Agric. Res. Serv., PSWCL Prog. Rep. 93-009.

Webster, J.A., C.A. Baker, and R.L. Burton. 1989. Progress in locating Russian wheat aphid resistance in barley and rye. Proc. Russian Wheat Aphid Conf., 25-27 October 1989, Albuquerque, NM, N. Mex. State Univ. Coop. Ext. Serv., pp. 18-19.

Webster, J.A., C.A. Baker, and D.R. Porter. 1991. Detection and mechanisms of Russian wheat aphid (Homoptera: Aphididae) resistance in barley. J. Econ. Entomol. 84:669-673.

Webster, J.A., and R.L. Burton. 1988. Host plant resistance studies and other research, pp. 14-19. In S. Coppock and B. Massey (eds.) Proceedings of the First Russian Wheat Aphid Conference. Okla. State Univ. Coop. Ext. Serv. Bull. E-875.

Webster, J.A., and R.L. Burton. 1989. The Russian wheat aphid: Its status and research on management strategies. Proc. 27th Barley Imp. Conf., pp. 57-61.

Webster, J.A., R.L. Burton, and K.J. Starks. 1987. *Diuraphis noxia*, a new U.S. aphid pests of small grains. J. Kans. Entomol. Soc. 60:483.

Webster, J.A., F. DuToit, and T.W. Popham. 1992. Reproduction of Russian wheat aphids in South Africa and Oklahoma, p. 210. In W.P. Morrison (comp.) Proceedings of the Fifth Russian Wheat Aphid Conference. Great Plains Agric. Counc. Pub. 142.



Webster, J.A., F. Dutoit, and T.W. Popham. 1993. Fecundity comparisons of the Russian wheat aphid (Homoptera: Aphididae) in Bethlehem, South Africa, and in Stillwater, Oklahoma. *J. Econ. Entomol.* 86:544-548.

Webster, J.A., C. Inayatullah, M. Hamissou, and K.A. Mirkes. 1994. Leaf pubescence effects in wheat on yellow sugarcane aphids and greenbugs (Homoptera: Aphididae). *J. Econ. Entomol.* 87:231-240.

Webster, J.A., and O.G. Merkle. 1988. Evaluating small grain germplasm for resistance to the Russian wheat aphid. *Proc. Beltsville Symp. in Agric. Res. XIII Biotic Diversity and Germplasm Preservation - Global Imperatives, Sec. II, No. 36.*

Webster, J.A., O.G. Merkle, and R.L. Burton. 1988. Russian wheat aphid plant resistance research in Oklahoma, pp. 46-47. *In* S. Coppock and B. Massey (eds.) *Proceedings of the First Russian Wheat Aphid Conference.* Okla. State Univ. Coop. Ext. Serv. Bull. E-875.

Webster, J.A., O.G. Merkle, and R.L. Burton. 1988. Progress in Russian wheat aphid plant resistance research, pp. 10-13. *In* F.B. Peairs and S.D. Pilcher (comp.) *Proceedings, 2nd Russian Wheat Aphid Workshop, 11-12 October 1988, Denver, CO.*

Webster, J.A., D.R. Porter, C.A. Baker, and D.W. Mornhinweg. 1993. Russian wheat aphid (Homoptera: Aphididae) resistance in barley: Effects on aphid feeding. *J. Econ. Entomol.* 86: 1603-1608.

Webster, J.A., and K.J. Starks. 1987. Fecundity of *Schizaphis graminum* and *Diuraphis noxia* (Homoptera: Aphididae) at three temperature regimes. *J. Kans. Entomol. Soc.* 60:580-582.

Webster, J.A., K.J. Starks, and R.L. Burton. 1987. Plant resistance studies with *Diuraphis noxia* (Homoptera: Aphididae), a new United States wheat pest. *J. Econ. Entomol.* 80:944-949.

Wright, S.P., T.J. Poprawski, W.L. Meyer, and F.B. Peairs. 1992. Natural prevalence of fungal pathogens, hymenopteran parasites, and dipteran predators of *Diuraphis noxia* and associated cereal aphid species on spring-planted grains in Colorado, p. 170-179. *In* W.P. Morrison (comp.) *Proceedings of the Fifth Russian Wheat Aphid Conference.* Great Plains Agric. Pub. 142.

Wright, S.P., T.J. Poprawski, W.L. Meyer, and F.B. Peairs. 1993. Natural enemies of the Russian wheat aphid (Homoptera: Aphididae) and associated cereal aphid species in spring-planted wheat and barley in Colorado. *Environ. Entomol.* 22:1383-1391.





